

Prevention of Football Injuries

Nader Rahnama¹

¹ MSc, MPhil, PhD, FLT, Associate Professor of Sport Injuries, School of Physical Education and Sports Sciences, University of Isfahan, Isfahan, Iran. Adjunct Professor of Sports Medicine, Department of Nutrition, Food and Exercise Sciences, Florida State University, Tallahassee, FL, USA.

Date of Submission: Dec 2, 2010

Date of Acceptance: Dec 25, 2010

Int J Prev Med 2011; 2(1): 38-40

Football (soccer) is certainly one of the most popular sports worldwide. It has been reported that more than 200,000 professional and 240 million amateur players play football.¹ Compared with other sports, football is a vigorous sporting activity with relatively high incidence of injury.²⁻⁵ To decrease the number of injuries, prevent early retirement, and provide a healthy and safe environment for players, preventive programs are highly recommended. For designing preventive programs, information about the incidence and risk of injury are required.⁵

The incidence, pattern and characteristics of football injuries such as injury type and nature, anatomical sites of injury, severity, causes, and mechanisms and also biomechanical, anthropometrical and physiological aspects of football players have been investigated and described in previous studies.⁶⁻³⁰ For example, a mean of 17-24 injuries per 1000 playing hours have been reported for football players²⁻⁴ and every male player generally encountered about one injury each year.^{30,31} Most of the epidemiological studies pointed out that the number of injuries during competition is about 4-6 times more than that in training.^{1,32,33} Sprains of the ankle and knee and strains of the hamstring and groin muscles are known as the four dominating injury types and account for more than 50% of all football injuries.³⁴ Over 65% of football injuries are classified as minor, 25% moderate, and 10% serious.³⁵ Approximately 50% of the football injuries occur from direct player-to-player contact, including collisions, tackling and being tackled, while the rest (non-contact) happen during the performance of actions such as running, shooting, turning, and heading.^{36,5} While

most investigators have illustrated that the occurrence of injury did not influenced by playing position,^{2,3} Hawkins and Fuller³⁶ and Zarei et al²³ reported that defenders had a greater risk of injury than other playing positions. Most injuries occur in the final 15 minutes of each half, and the risk of injury in the second half is greater than the first half.³⁶⁻³⁸

Prevention of injury would maintain the health of player, minimize costs, and also enhance performance.³⁹ Only a few studies have been conducted on football injury prevention. Engebretsen and Bahr⁴⁰ reported that out of 10,691 papers published on athletic injury, there were only 6 randomized controlled trials on sports injury prevention. Regarding the published papers on prevention of injury in football in the last three decades, no more than 15 papers have been published on injury prevention. The first injury prevention study was conducted by Ekstrand et al 28 years ago.⁴¹ They found a significant reduction in the number of injuries through a 7-part prevention program. Two years later, Tropp et al⁴² following a balance training program observed a significantly lower number of ankle sprains in football players. The positive preventive effect of orthoses, proprioceptive and eccentric strength training programs on prevention of ankle, knee and hamstring injuries were reported.⁴³⁻⁴⁶ Caraffa et al⁴³ found a lower incidence of ACL injuries in semi-professional and amateur football players following a special proprioceptive training program. Heidt et al⁴⁷ planned an injury prevention program (pre-season conditioning program) in female players and observed lower incidence of injury (14%) compare to the untrained group. Junge et al⁴⁸

showed 21% fewer injuries in male youth football players following an interventional programs educating and supervising players and coaches.

Consequently, injury prevention programs in football will definitely need to be developed. By conducting more studies about the risk factors and associated roles, researchers will be able to provide sound advices for players, team medical staff, coach, and referees may help them to make a safe environment for football players. It seems that the co-operation of important governmental and international organizations such as FIFA, UEFA, FA, IOC and special football federations of each country is vital. Studies on the effectiveness of injury prevention and interventional programs should be considered and expected. Since information on effectiveness of intervention and also data on extrinsic and intrinsic risk factors in football are very beneficial for minimizing injury, they should be top priorities for injury prevention studies and because of limitation of resources for injury prevention studies, it is suggested that effective and comprehensive studies be designated and funded.⁴⁹ Further studies, preferably randomized controlled trails of the effectiveness of football injury prevention interventional studies should be undertaken as they provide the most valuable and valid data. Finally, too many conferences in the field of sports medicine, especially on prevention of football injury should be established and organized worldwide.

Conflict of interest statement: The author declares that he has no conflict of interest.

Source of funding: None

REFERENCES

1. Junge A, Dvorak J. Soccer injuries: a review on incidence and prevention. *Sports Med* 2004; 34(13): 929-38.
2. Engstrom B, Johansson C, Tornkvist H. Soccer injuries among elite female players. *Am J Sports Med* 1991; 19(4): 372-5.
3. Hawkins RD, Fuller CW. An examination of the frequency and severity of injuries and incidents at three levels of professional football. *Br J Sports Med* 1998; 32(4): 326-32.
4. Hawkins RD, Fuller CW. A prospective epidemiological study of injuries in four English professional clubs. *Br J Sports Med* 1999; 33(3): 196-203.
5. Rahnama N, Reilly T, Lees A. Injury risk associated with playing actions during competitive soccer. *Br J Sports Med* 2002; 36(5): 354-9.
6. Rahnama N, Lees A, Reilly T. Electromyography of selected lower-limb muscles fatigued by exercise at the intensity of soccer match-play. *J Electromyogr Kinesiol* 2006; 16(3): 257-63.
7. Rahnama N, Manning LK. The mechanisms and characteristics of injuries in English youth-soccer. In: Reilly T, Cabri J, Araujo D, Editors. *Science and football V: the proceedings of the Fifth World Congress on Science and Football*. London: Routledge; 2005. p. 302-6.
8. Rahnama N, Reilly T, Lees A, Graham-Smith P. Muscle fatigue induced by exercise simulating the work rate of competitive soccer. *J Sports Sci* 2003; 21(11): 933-42.
9. Rahnama N, Reilly T, Lees A. A novel computerised notation and analysis system for assessment of injury and injury risk in football. *Phys Ther Sport* 2002; 3(4): 183-90.
10. Rahnama N, Zareei M. Video analysis of football injuries at the Asian Cup 2007. In: Drust B, Reilly T, Williams AM, Editors. *International Research in Science and Soccer*. London: Routledge; 2009. p. 123-31.
11. Rahnama N, Lees A, Bambaecichi E. Comparison of muscle strength and flexibility between the preferred and non-preferred leg in English soccer players. *Ergonomics* 2005; 48(11-14): 1568-75.
12. Lees A, Steward I, Rahnama N, Barton G. Lower limb function in the maximal instep kick in soccer. In: Reilly T, Atkinson G, Editors. *Contemporary Sport, Leisure and Ergonomics*. New York: Taylor & Francis; 2009. p. 149-60.
13. Rahnama N, Reilly T, Lees A. An analysis of injury risk in the Manchester united team when playing at home and away. *Cellular & Molecular Biology Letters* 2004; 9(4): 773-6.
14. Tarmah H, Rahnama N, Khayyambashi K. Video analysis of causes and mechanisms of the ACL injuries in the Iranian professional soccer players. *Br J Sports Med* 2010; 44(Suppl 1): 3-4.
15. Zarei M, Rahnama N, Rezaei J, Reisi J. Comparison of Injury in Iran national junior and youth football players. *Br J Sports Med* 2010; 44(Suppl 1): 6.
16. Rahnama N, Reilly T, Lees A, Graham-Smith P. A comparison of musculoskeletal function in elite and sub-elite English soccer players. In: Reilly T, Marfell-Jones M, Editors. *Kinanthropometry VIII: proceedings of the 8th International Conference of the International Society for the Advancement of Kinanthropometry (ISAK)*. London: Routledge; 2003. p. 151-64.
17. Zarei M, Rahnama N, Bambaecichi E. Performance and injury incidence rate in home and away games in Iran premier league teams. *Br J Sports Med* 2010; 44(Suppl 1): 15.
18. Nazarian AB, Khayyambashi K, Rahnama N, Salamat MR. The comparison of bone mineral density in lumbar spines and femoral bone between professional soccer players and non-athlete subjects. *WJSS* 2009; 2(2): 106-11.

19. Rahnema N, Reilly T, Lees A. Electromyographic analysis of selected muscles during a simulated soccer game. In: Reilly T, Cabri J, Araujo D, Editors. *Science and football V: the proceedings of the Fifth World Congress on Science and Football*. London: Routledge; 2005. p. 509-15.
20. Nazarian AB, Khayambashi K, Rahnema N. Dominant and non-dominant leg bone mineral density in professional soccer players and non-athlete subjects. *WJSS* 2010; 3(1): 28-32.
21. Rahnema N, Bambaiechi E, Daneshjoo H. The epidemiology of knee injuries in Iranian male professional soccer players. *Sport Sci Health* 2009; 5(1): 9-14.
22. Rahnema N, Younesian A, Mohammadion M, Bambaiechi E. Effect of 90-min soccer match on lipid profiles in professional players. *JRMS* 2009; 14(6): 335-41.
23. Zarei M, Rahnema N, Rajabi R. The effect of soccer players' positional role in Iran super league on sport injury rates. *WJSS* 2009; 2(1): 60-4.
24. Lees A, Steward I, Rahnema N, Barton G. Understanding lower limb function in the maximal instep kick in soccer. *J Sport Sci* 2009; 27: S20-S21.
25. Rahnema N, Naderian M, Ebrahimi E, Bambaiechi E. Incidence of injury between guest and host football teams. *Journal of Sports Science & Medicine* 2007; 6(Supplementary 10): S24.
26. Rahnema N, Bambaiechi E, Sadeghipour HR. Home advantage in Asian football leagues. *Journal of Sports Science & Medicine* 2007; 6(Suppl 10): S209.
27. Sajjadi N, Rahnema N. Analysis of goals in 2006 FIFA World Cup. *Journal of Sports Science & Medicine* 2007; 6(Suppl 10): S3.
28. Rahnema N, Lees A, Barton G. The error in kinetic data for the maximal soccer instep kick associated with standing calibration. *Journal of Sports Sciences* 2005; 23: S110.
29. Lees A, Rahnema N, Barton G. The stability and between-day reliability of kinetic data for the maximal soccer instep kick associated with standing calibration. *Journal of Sports Sciences* 2005; 23: 110-1.
30. Dvorak J, Junge A. Football injuries and physical symptoms. A review of the literature. *Am J Sports Med* 2000; 28(5 Suppl): S3-S9.
31. Hawkins R, Hulse M, Wilkinson C, Hodson A, Gibson M. The association football medical research programme: an audit of injuries in professional football. *Br J Sports Med* 2001; 35(1): 43-7.
32. Lewin G. The incidence of injury in an English professional soccer club during one competitive season. *Physiotherapy* 1989; 75(10): 601-5.
33. McGregor JC, Rae A. A review of injuries to professional footballers in a premier football team (1990-93). *Scott Med J* 1995; 40(1): 16-18.
34. Engebretsen AH, Myklebust G, Holme I, Engebretsen L, Bahr R. Prevention of injuries among male soccer players: a prospective, randomized intervention study targeting players with previous injuries or reduced function. *Am J Sports Med* 2008; 36(6): 1052-60.
35. Anglietti P, Zaccherotti G, De Biase P. Injuries in soccer: mechanism and epidemiology. In: Renstrom P, editor. *Clinical practice of sports injury prevention and care*. New York: Wiley-Blackwell; 1994. p. 277-84.
36. Hawkins RD, Fuller CW. Risk assessment in professional football: an examination of accidents and incidents in the 1994 World Cup finals. *Br J Sports Med* 1996; 30(2): 165-70.
37. Sandelin J, Santavirta S, Kiviluoto O. Acute soccer injuries in Finland in 1980. *Br J Sports Med* 1985; 19(1): 30-3.
38. Arnason A, Gudmundsson A, Dahl HA, Johannsson E. Soccer injuries in Iceland. *Scand J Med Sci Sports* 1996; 6(1): 40-5.
39. Meeuwisse W, Bahr R. A systematic approach to sports injury prevention. In: Bahr R, Engebretsen L, Editors. *Sports Injury Prevention*. New Jersey: John Wiley and Sons; 2009. p. 7-16.
40. Engebretsen L, Bahr R. Why injury prevention in sports important? In: Bahr R, Engebretsen L, Editors. *Sports Injury Prevention*. New Jersey: John Wiley and Sons; 2009. p. 1-6.
41. Ekstrand J, Gillquist J, Liljedahl SO. Prevention of soccer injuries. Supervision by doctor and physiotherapist. *Am J Sports Med* 1983; 11(3): 116-20.
42. Tropp H, Askling C, Gillquist J. Prevention of ankle sprains. *Am J Sports Med* 1985; 13(4): 259-62.
43. Caraffa A, Cerulli G, Proietti M, Aisa G, Rizzo A. Prevention of anterior cruciate ligament injuries in soccer. A prospective controlled study of proprioceptive training. *Knee Surg Sports Traumatol Arthrosc* 1996; 4(1): 19-21.
44. Surve I, Schwellnus MP, Noakes T, Lombard C. A fivefold reduction in the incidence of recurrent ankle sprains in soccer players using the Sport-Stirrup orthosis. *Am J Sports Med* 1994; 22(5): 601-6.
45. Askling C, Karlsson J, Thorstensson A. Hamstring injury occurrence in elite soccer players after preseason strength training with eccentric overload. *Scand J Med Sci Sports* 2003; 13(4): 244-250.
46. Arnason A, Andersen TE, Holme I, Engebretsen L, Bahr R. Prevention of hamstring strains in elite soccer: an intervention study. *Scand J Med Sci Sports* 2008; 18(1): 40-8.
47. Heidt RS, Jr., Sweeterman LM, Carlonas RL, Traub JA, Tekulve FX. Avoidance of soccer injuries with preseason conditioning. *Am J Sports Med* 2000; 28(5): 659-62.
48. Junge A, Rosch D, Peterson L, Graf-Baumann T, Dvorak J. Prevention of soccer injuries: a prospective intervention study in youth amateur players. *Am J Sports Med* 2002; 30(5): 652-9.
49. Roberts I. Research priorities for injury prevention. *Inj Prev* 2001; 7(1): 2-3.